

STARTING THE CONVERSATION ABOUT A *once-monthly** TREATMENT OPTION



*You should continue to take your current antipsychotic medication for 2 weeks after your first treatment with ABILIFY MAINTENA.



Here are some questions that can make having a discussion with your loved one's healthcare provider easier—and help in determining whether once-monthly* ABILIFY MAINTENA® (aripiprazole) is right for your loved one.

- How do you recommend helping my loved one understand their treatment options? _____
- Could a once-monthly treatment like ABILIFY MAINTENA be an appropriate option for my loved one? _____
- What are the possible side effects of ABILIFY MAINTENA? _____
- Do my loved one's schizophrenia symptoms need to be under control before starting ABILIFY MAINTENA? _____
- Does my loved one need to stay on their current treatment after starting ABILIFY MAINTENA? _____
- How is ABILIFY MAINTENA given? _____
- What are the potential advantages and disadvantages of a long-acting injectable treatment? _____
- How could ABILIFY MAINTENA help my loved one work towards their treatment goals? _____

INDICATIONS and IMPORTANT SAFETY INFORMATION for ABILIFY MAINTENA® (aripiprazole)

INDICATIONS:

ABILIFY MAINTENA is a prescription medicine given by injection by a healthcare professional:

- for the treatment of schizophrenia in adults
- alone as maintenance treatment of bipolar I disorder in adults

It is not known if ABILIFY MAINTENA is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION:

Increased risk of death in elderly people with dementia-related psychosis. ABILIFY MAINTENA increases the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). ABILIFY MAINTENA is not for the treatment of people with dementia-related psychosis.

(continued on next page)

Please read [FULL PRESCRIBING INFORMATION](#), including **BOXED WARNING**, and [MEDICATION GUIDE](#).

IMPORTANT SAFETY INFORMATION FOR ABILIFY MAINTENA® (aripiprazole) (cont'd)

Do not receive ABILIFY MAINTENA if you are allergic to aripiprazole or any of the ingredients in ABILIFY MAINTENA.

ABILIFY MAINTENA may cause serious side effects, including:

- **Increased risk of stroke** and ministroke has been reported in clinical studies with oral aripiprazole of elderly people with dementia-related psychosis and can lead to death.
- **Neuroleptic malignant syndrome (NMS), a serious condition that can lead to death.** Call your healthcare provider or go to the nearest emergency room right away if you have some or all of the following symptoms of NMS: high fever, stiff muscles, confusion, sweating, or changes in pulse, heart rate, and blood pressure.
- **Uncontrolled body movements (tardive dyskinesia).** ABILIFY MAINTENA may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop receiving ABILIFY MAINTENA. Tardive dyskinesia may also start after you stop receiving ABILIFY MAINTENA.
- **Problems with your metabolism such as:**
 - » **High blood sugar (hyperglycemia) and diabetes:** Increases in blood sugar can happen in some people who receive ABILIFY MAINTENA. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes, such as being overweight or a family history of diabetes, your healthcare provider should check your blood sugar before you start receiving ABILIFY MAINTENA and during your treatment.

Call your healthcare provider if you have any of these symptoms of high blood sugar while receiving ABILIFY MAINTENA:

- feel very thirsty
- feel weak or tired
- need to urinate more than usual
- feel sick to your stomach
- feel very hungry
- feel confused, or your breath smells fruity
- » **Increased fat levels (cholesterol and triglycerides) in your blood.**
- » **Weight gain.** You and your healthcare provider should check your weight regularly during treatment.
- **Unusual and uncontrollable (compulsive) urges.** Some people receiving ABILIFY MAINTENA have had unusual strong urges to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having unusual urges or behaviors, talk to your healthcare provider.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** ABILIFY MAINTENA may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills, which may lead to falls that can cause fractures or other injuries.
- **Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment.
- **Seizures (convulsions)**
- **Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities.** Do not drive a car, operate machinery, or do other dangerous activities until you know how ABILIFY MAINTENA affects you. ABILIFY MAINTENA may make you feel drowsy and affect your judgement, thinking, or motor skills.

- **Problems controlling your body temperature so that you feel too warm.** Do not become too hot or dehydrated while you receive ABILIFY MAINTENA. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing** that can cause food or liquid to get into your lungs.

Do not drink alcohol while you receive ABILIFY MAINTENA.

Before receiving ABILIFY MAINTENA, tell your healthcare provider about all of your medical conditions, including if you:

- have never taken aripiprazole before
- have or had diabetes or high blood sugar or a family history of diabetes or high blood sugar
- have or had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or had low or high blood pressure
- have or had heart problems or a stroke
- have or had a low white blood cell count
- have or had seizures (convulsions)
- have problems that may affect you receiving an injection in your arm or buttocks
- are pregnant or plan to become pregnant. ABILIFY MAINTENA may harm your unborn baby. If you become pregnant during treatment with ABILIFY MAINTENA, talk to your healthcare provider about the risk to your unborn baby and about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>
- are breastfeeding or plan to breastfeed. ABILIFY MAINTENA can pass into your milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you receive ABILIFY MAINTENA.

Tell your healthcare provider about all the medicines you take, including prescription medicines and over-the-counter medicines, vitamins, and herbal supplements.

ABILIFY MAINTENA and other medicines may affect each other causing possible serious side effects. Do not start or stop any medicines during treatment with ABILIFY MAINTENA without talking to your healthcare provider first.

The most common side effects of ABILIFY MAINTENA include: weight gain, restlessness or feeling like you need to move (akathisia), injection site pain, or sleepiness (sedation).

It is important to contact your healthcare provider if you experience prolonged, abnormal muscle spasms or contractions, which may be signs of a condition called dystonia.

These are not all the possible side effects of ABILIFY MAINTENA.

If you have any questions about your health or medicines, talk to your healthcare provider.

You are encouraged to report side effects of ABILIFY MAINTENA. Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please see [FULL PRESCRIBING INFORMATION](#), including **BOXED WARNING**, and [MEDICATION GUIDE](#) for ABILIFY MAINTENA.

